

# CHEMICAL COMPOSITION AND ANTIMICROBIAL ACTIVITY OF THE ESSENTIAL OIL FROM *HERACLEUM SIAMICUM*

Tiwatt Kuljanabhadgavad<sup>1,3</sup>, Nongluksna Sriubolmas<sup>2</sup> and Nijisiri Ruangrungsi<sup>1,\*</sup>

<sup>1</sup>Department of Pharmacognosy and Pharmaceutical Botany,

<sup>2</sup>Department of Biochemistry and Microbiology, Faculty of Pharmaceutical Sciences, Chulalongkorn University, Bangkok 10330, Thailand, <sup>3</sup>Bioorganic Chemistry Research Laboratories, Faculty of Science and Technology, Suan Dusit Rajabhat University, Bangplad, Bangkok 10700, Thailand

**ABSTRACT:** *Heracleum siamicum* Craib (Apiaceae) is an important herbal spices having a wide applications in flavoring processed foods. The flat-oval shaped fruit of *H. siamicum* Craib from North Thailand was hydrodistilled and chemical composition of the essential oil was analyzed by GC and GC-MS. The essential oil yield based on dried plant material was 1.25% and twenty-five compounds (corresponding to 97.69% of the total weight) were identified. The main components were: *n*-octyl acetate (65.30%), *o*-cymene (10.35%), limonene (7.52%),  $\delta$ -2-carene (6.87%), *cis*-thujone (1.92%), isobornyl acetate (0.94%), *n*-octanol (0.73%), 1,8-cineol (0.62%), *n*-tridecanol (0.44%), and safrole (0.37%). *H. siamicum* essential oil demonstrated bactericidal and fungicidal activity against five bacterial strains and two fungal strains, using agar diffusion and minimum inhibitory concentration methods.

**Keywords:** *Heracleum siamicum*, antimicrobial activity, chemical composition, hydrodistillation, volatile oil analysis, GC-MS

**INTRODUCTION:** *Heracleum siamicum* Craib (Apiaceae) is a perennial sturdy plant known as “Ma Laep” found in the northern (N) and northeast (NE) parts of Thailand<sup>1</sup>. The fruits of *H. siamicum* are widely used as spices. In Thai folk medicine, the fruits of *H. siamicum* were used as a carminative herbal drug. Because of wide usage of the fruits of *H. siamicum* as medicinal plant and its use as flavoring agent, it was decided to carry out a phytochemical study on the fruit of this plant.

Many kinds of metabolites including coumarins, furanocoumarins, anthraquinones, stilbenes, furanocoumarin dimers, and flavonoids have been isolated and identified from various species of this genus<sup>2-10</sup>. Plants belonging to the genus *Heracleum* are aromatic and are excellent sources of essential oils. The essential oil composition of various members of this genus have been reported, *H. persicum*<sup>11,12</sup>, *H. candolleianum* Wight et Arn. Gamble<sup>13</sup>, *H. dissectum* Ledeb.<sup>14</sup>, *H. sphondylium* L. subsp. *ternatum* (Velen.) Brummitt<sup>15</sup>, *H. crenatifolium* Boiss<sup>16, 17</sup>, *H. platytaenium* Boiss.<sup>16</sup>, and *H. candolleianum*<sup>18</sup> contain monoterpene hydrocarbons (e.g. *p*-cymene;  $\gamma$ -terpene;  $\alpha$ - and  $\beta$ -pinene; limonene etc.), oxygenated monoterpenes (e.g. iso-bornyl acetate, linalool, *n*-octanol, terpinene-1-ol-4 etc.), and sesquiterpene (e.g.

caryophyllene oxide) in their volatile fractions. The extracts from the fruits of *H. persicum* Desf. ex Fisher showed antibacterial activity that inhibited the growth of *Staphylococcus aureus*, *Bacillus cereus*, and *Bordetella bronchiseptica*<sup>19</sup>. Different octyl esters, especially *n*-octyl acetate, is reported to be the major constitute in most of the oils investigated<sup>12,16,17,20,21</sup>. The volatile substances from *H. siamicum* fruits and their antimicrobial activity are reported for the first time in this communication. This study concerns chemical composition and antimicrobial activity of the essential oil from *H. siamicum*.

## MATERIALS AND METHODS:

### *Plant material and Isolation of the Essential Oil*

Fruit of *H. siamicum* Craib, was collected in January 2008 from the market of Chiangmai Province in the north of Thailand. A voucher specimen was deposited in the Department of Pharmacognosy and Pharmaceutical Botany, Chulalongkorn University. The dried fruits were hydrodistilled in a Clevenger-type apparatus, according to the literature<sup>22</sup>. The oil was dried over anhydrous sodium sulfate and stored at 4°C in a vial covered with aluminum foil to prevent the negative effect of light and submitted to chemical and microbiological analysis.

\*To whom correspondence should be addressed.  
E-mail: nijisiri.r@chula.ac.th, Tel. +668 1917 9541

### Analysis of Essential Compound

Analysis was performed with a Varian Star 3400 CX gas chromatograph coupled with a Saturn III mass spectrometer (Varian Inc.) system equipped with a Varian automatic injector and a 30 m long, DB-5 MS (J&W) capillary column (0.25 mm id, 0.25  $\mu$ m film thickness). The ionization energy was 70 eV. A sample of 1.0  $\mu$ l of a 4% solution of the fruit oil in hexane was injected with a split ratio of 100:1. The temperature of the injection block was 240°C. The GC oven temperature was programmed as follows: initial temperature 60°C (1 min) followed by a temperature increase of 3°C/min up to 200°C and a second ramp of 5°C/min to the final temperature of 220°C. The carrier gas was helium at 1.0 ml/min at constant volume. Identification of the oil components was established by comparing GC-MS spectra and RI with those of an internal Varian NIST MS 1998 library and those described by Adams<sup>23</sup>.

### Antimicrobial Activity

The microbial strains used in the antimicrobial assays were: the gram-positive bacteria *Bacillus subtilis* (ATCC 6633), *Staphylococcus aureus* (ATCC 29213), *Streptococcus faecalis* (ATCC 29212) and the gram-negative bacteria *Escherichia coli* (ATCC 25922), *Pseudomonas aeruginosa* (ATCC 27853) and the pathogenic fungi *Candida albicans* (ATCC 10231) and *Microsporium gypseum* (a clinical isolate). Antimicrobial activities of the volatile oil of *H. siamicum* were determined using the agar-disc diffusion method<sup>24-28</sup>, as described below. Each bacterial strain was grown on Trypticase Soy Agar plates at 37°C for 24 h. Portions of four discrete colonies were inoculated into 5 ml of Trypticase Soy Broth (TSB) and incubated at 37°C for 2–3 h. The turbidity of each culture was adjusted with sterile saline. For yeast, *C. albicans* was grown on Sabouraud Dextrose Agar (SDA) slant at 30°C for 24 h and some of the growth was transferred to 5 ml of sterile saline. Turbidity of the inoculum suspension was adjusted with sterile saline. *Microsporium gypseum* (of the mold spore) was grown on SDA at 30°C for 96 h, washed from the slant culture and adjusted

to the desired turbidity with sterile 0.05% Tween 80. Additionally, the plates with internal diameter of 100 mm containing 25 ml of Muller-Hinton agar and SDA were inoculated with bacterial suspension and fungal suspension by streaking method respectively<sup>25</sup>. The wells (6 mm holes) were produced in the agar with sterile cork borer No.3. The fruit oils were diluted with sterile 0.05% Tween 80 to the final concentration of 1:20. Of this, 50  $\mu$ L of the diluted samples were pipetted into each well. The plates were left at room temperature for 1 h and then incubated at 37°C for 24 h for bacteria and at 30°C for 96 h for fungi. The tests were carried out in duplicate. The minimum inhibitory concentrations of the oil using the dilution assays were determined as previously described<sup>26</sup>. Reading the results was carried out by measuring the diameters of the zones of inhibition and clear growth (in mm) and the minimum inhibitory concentration (MIC) was defined as the lowest concentration of the volatile oil which prevented growth of the inoculum compared with the growth control plate.


**RESULTS AND DISCUSSION:** Fruits contained 1.25% (v/w) essential oil (dried weight) was light yellow liquid in color and possessed a distinct sharp odor. Table 1 and Figure 1 showed its chemical composition. Twenty-five compounds were identified by comparison of their retention indexes and the mass spectra of each GC component with those of standards and with reported data. Terpenes and their derivatives predominated, with the most abundant one being *n*-octyl acetate (65.30%), followed by *o*-cymene (10.35%), limonene (7.52%),  $\delta$ -2-carene (6.87%), *cis*-thujone (1.92%), isobornyl acetate (0.94%), *n*-octanol (0.73%), 1,8-cineol (0.62%), *n*-tridecanol (0.44%), and safrole (0.37%), respectively. These main components comprised more than 95% of the essential oil. We should also note the presence in the essential oil of a total 4.31 % of alcohol hydrocarbons (Table 2). Although most of these compounds are well documented as essential oil components in various plant species<sup>29</sup>, to our knowledge this is the first reported of their occurrence in the essential oil of *H. siamicum*.

**Table 1** Chemical composition of essential oil from *Heracleum siamicum* determined by GC-MS

No	Compound	RI*	Percent
1	Tricyclene	926	0.22 <sup>a, c</sup>
2	$\alpha$ -Thujene	930	0.21 <sup>a</sup>
3	$\delta$ -2-Carene	1001	6.87 <sup>a, c</sup>
4	$\delta$ -3-Carene	1011	0.33 <sup>a, c</sup>
5	$\alpha$ -Terpinene	1017	0.07 <sup>a</sup>
6	<i>o</i> -Cymene	1022	10.35 <sup>b</sup>
7	Limonene	1030	7.52 <sup>b, c</sup>
8	1,8-Cineol	1033	0.62 <sup>a, c</sup>
9	<i>n</i> -Octanol	1070	0.73 <sup>a</sup>
10	Linalool	1097	0.13 <sup>b</sup>
11	<i>cis</i> -Thujone	1102	1.92 <sup>a</sup>
12	<i>trans</i> -Pinocarveol	1139	0.10 <sup>a, b</sup>
13	Camphor	1143	0.26 <sup>b, c</sup>
14	Borneol	1165	0.10 <sup>b, c</sup>
15	Terpin-4-ol	1177	0.13 <sup>a, b</sup>
16	<i>n</i> -Octyl acetate	1194	65.30 <sup>b</sup>
17	Isobornyl acetate	1285	0.94 <sup>a</sup>
18	Safrole	1285	0.37 <sup>a, c</sup>
19	$\alpha$ -Copaene	1376	0.36 <sup>a</sup>
20	$\beta$ -Bourbonene	1384	0.27 <sup>a</sup>
21	9- <i>epi</i> -( <i>E</i> )-Caryophyllene	1467	0.14 <sup>a, c</sup>
22	Citronellyl isobutyrate	1482	0.11 <sup>a</sup>
23	Viriflorene	1493	0.39 <sup>a</sup>
24	$\delta$ -Cadinene	1523	0.24 <sup>a</sup>
25	<i>n</i> -Tridecanol	1575	0.44 <sup>a, c</sup>

\*RI determined on a DB-5 column using the homologous series of *n*-hydrocarbons (Kovats index).

<sup>a</sup>Identification was based on RI. <sup>b</sup>Identification was based on comparison of the GC-MS spectra and RI with those of internal (computer) NIST library and those described by Adams. <sup>c</sup>Identification was based on comparison of authentic standards.

**Figure 1** Structure of the major components of essential oil of *H. siamicum*

The dominant compound, *n*-octyl acetate, has been reported as a common component in most fruit oils of *Heracleum* genus and also reported as the constitute in *Boswellia carterii* Birdw<sup>30</sup>, *Peucedanum cervaria* (L.) Lapeyr<sup>31</sup>, and grapefruit oil<sup>32</sup>. In a recently, the essential oil of *H. sphondylium* subsp. *ternatum* contain the major constitute as *n*-octanol had a high antimicrobial activity against *Candida albicans* with a MIC of 0.5 mg/ml<sup>15</sup>. Limonene has been shown to be biologically active as an antitumour agent<sup>33</sup>. The essential oil of *Grammosciadium platycarpum* Boiss. contained the major constitute as limonene had a high antimicrobial activity against *Staphylococcus epidermidis* with a MIC of 0.6–5 mg/ml<sup>34</sup>.

Interestingly, there were significant differences between the main components of the essential oil of *H. siamicum* Craib and those previously determined in *H. crenatifolium* Boiss.<sup>17</sup> which belongs to the same genus. Thus, terpene alcohols such as *n*-octanol, limonene, and linalool are quantitatively abundant in *H. candolleianum* oil, whilst they were only present in much smaller quantities in *H. siamicum* oil (Table 3).

Results of the antimicrobial activity tests of *H. siamicum* essential oil was examined by agar disc diffusion assay and minimum inhibition concentration against an array of five bacteria and two fungi selected on the basis of their relevance to public health (Table 4). The oil demonstrated strong bacteriostatic rather than fungistatic activities. Particularly significant were the inhibition zone diameters observed for dilution essential oil against *Staphylococcus aureus*, *Bacillus subtilis*, and *Candida albicans*, while *Escherichia coli*, *Pseudomonas aeruginosa* and *Microsporium gypseum* were less sensitive to the oil.

The oil showed an MIC of 20  $\mu$ g/ml against *S. aureus*, 25  $\mu$ g/ml against *B. subtilis*, and 50  $\mu$ g/ml against *C. albicans*, respectively. Moreover, the paper clearly suggests that the chemical composition ( $\delta$ -2-carene, *o*-cymene, *cis*-thujone, limonene, isobornyl acetate, *n*-octanol, 1,8-cineol, linalool, and safrole) show synergistic effect which is much more greater than the sum of anti-bacterial effect of each component used alone<sup>35-39</sup>.

**Table 2** Composition of *H. siamicum* essential oil by substance classes

Compounds	% in essential oil
Monoterpenes	25.14
Sesquiterpenes	1.40
Saturated	66.47
<b>Hydrocarbon total:</b>	<b>93.01</b>
Alcohols	4.31
Deoxymethylene	0.37
<b>Oxygenated compounds total:</b>	<b>4.68</b>
<b>Total compounds:</b>	<b>97.69</b>

**Table 3** Main composition of the essential oils from *H. siamicum* and *H. crenatifolium*<sup>17)</sup>

<i>H. siamicum</i> (Relative amount, %)	<i>H. crenatifolium</i> (Relative amount, %)
<i>n</i> -octyl acetate (65.30%)	octyl acetate (88.4%)
<i>o</i> -cymene (10.35%)	octanol (3.10%)
limonene (7.52%)	( <i>Z</i> )-4-octenyl acetate (1.0%)
$\delta$ -2-carene (6.87%)	octyl 2-methyl butyrate (0.9%)
<i>cis</i> -thujone (1.92%)	octyl hexanoate (0.7%)
isobornyl acetate (0.94%)	hexyl 2-methyl butyrate (0.7%)
<i>n</i> -octanol (0.73%)	$\alpha$ -pinene (0.7%)
1,8-cineol (0.62%)	octanal (0.6%)
<i>n</i> -tridecanol (0.44%)	myristicin (0.4%)
safrole (0.37%)	limonene (0.3%)

**Table 4** Antimicrobial activity of essential oil from *H. siamicum*

Tested Microorganism*	Inhibition zone (mm) <sup>a</sup>	MIC ( $\mu$ g/ mL)
<i>Bacillus subtilis</i> ATCC 6633	11.23 $\pm$ 0.73	25
<i>Candida albicans</i> ATCC 10231	9.70 $\pm$ 0.93	50
<i>Escherichia coli</i> ATCC 25922	-	-
<i>Streptococcus faecalis</i> ATCC 29212	-	-
<i>Streptococcus aureus</i> ATCC 29213	11.43 $\pm$ 1.01	20
<i>Microsporium gypseum</i> (a clinical isolate)	-	-
<i>Pseudomonas aeruginosa</i> ATCC 27853	-	-

\*Tested in 50  $\mu$ l of 10% oil in Tween 80;

<sup>a</sup> mean values  $\pm$  SD, The 0.05% Tween 80 did not show any activity, - No inhibition zone.

In addition to the individual monoterpene alcohol of *Salvia fructicosa* Mill. (Lamiaceae), such as 1,8-cineole, and thujone, a major constituents showed relatively high levels of antimicrobial activity against the bacteria, while camphor was inactive against the bacteria tested<sup>33)</sup>. However, it is possible that the activity of the main components is modulated by other minor molecules<sup>39-41)</sup>. In general, the cytotoxic activity of

essential oils is mostly due to the presence of aldehydes, alcohols, methylene dioxy compounds, and phenols<sup>28,42-43)</sup>.

Our group has previously reported on the antimicrobial activity of *Cinnamomum porrectum* (Roxb.) Kosterm (collected from E. Thailand) essential oil that has safrole (99.8%) as the main component had a very high antimicrobial activity against *Candida albicans* with a MIC of 0.063% by volume<sup>29)</sup>. Moreover, *C. porrectum* oil provided a potential for treatment of *Candida* infections and prepared in solution and emulgel topical dosage forms at various concentration (1, 2, and 5% w/w)<sup>44)</sup>. *H. siamicum* has moderate oil content and a high proportion of monoterpenes in its profile. The dominance of *o*-cymene, limonene, and the presence of isobornyl acetate in the profile make *H. siamicum* a good candidate for further study in terms of its allelochemical properties, and the agreeable fragrance of the oil makes it of possible interest to the flavoring agent in food and in perfumery.

**CONCLUSION:** Our GC and GC-MS study of the essential oil of *H. siamicum* from N.Thailand led to the identification of twenty-five compounds, representing 97.69% of the total mass. The main constituents were terpenes and their derivatives, and the most prominent one was *o*-cymene (10.35%). The antimicrobial activity results presented here demonstrate that this plant essential has a commercial potential.

**ACKNOWLEDGMENTS:** This research was funded by the Biodiversity Research and Training Program (BRT) to NR. We are grateful to the Scientific and Technological Research Equipment Center, CU for the Varian GC and GC-MS instrument. The identification of the plant was confirmed by one of us (NR). Thanks to Professor Michael Wink, Institute of Pharmacy and Molecular Biotechnology (IPMB), University of Heidelberg, Germany for his comments and the reading of the manuscript.

#### REFERENCES:

- Hedge IC, Lamond JM. 1992. *Umbelliferae*. The Chutima Printing; Bangkok,; Vol. 5, p 442.
- Doi M, Nakamori T, Shibano M, Taniguchi M, Wang N-H, Baba K. 2004. Candibirin A, a

- furanocoumarin dimer isolated from *Heracleum candicans* Wall. Acta Crystallogr C 60: o833–5.
3. Razdan TK, Kachroo V, Harkar S, Koul GL. 1982. Furanocoumarins from *Heracleum canescens*. Phytochem 21(4): 923–7.
  4. Taniguchi M, Yokota O, Shibano M, Wang NH, Baba K. 2005. Four coumarins from *Heracleum yunnngingense*. Chem Pharm Bull 53(6): 701–4.
  5. Ghodsi B. 1976. Flavonoids of three *Heracleum* species: *H. Persicum* L., *H. sphondylium* L. and *H. montanum* Schl. Bull Trav Soc Pharm Lyon 20: 3–8.
  6. Bogucka-Kocka A, Rulka J, Kocki J, Kubis P, Buzala E. 2004. Bergapten apoptosis induction in blood lymphocytes of cattle infected with *bovine leukaemia virus* (BLV). Bull Vet Inst Pulawy 48(2): 99–103.
  7. Niu XM, Li SH, Wu LX, Li L, Gao LH, Sun HD. 2004. Two new coumarin derivatives from the roots of *Heracleum rapula*. Planta Med 70(6): 578–81.
  8. Ceska O, Chaudhary SK, Warrington PJ, Ashwoodsmith MJ. 1986. Photoactive furocoumarins in fruits of some umbellifers. Phytochem 26(1): 165–9.
  9. Orhan I, Tosun F, Şener B. 2008. Coumarin, anthroquinone and stilbene derivatives with anticholinesterase activity. Z Naturforsch 63c: 366–70.
  10. Sayyah M, Moaied S, Kamalinejad M. 2005. Anticonvulsant activity of *Heracleum persicum* seed. J Ethnopharmacol 98(1–2): 209–11.
  11. Sefidkon F, Dabiri M, Mohammad N. 2004. Analysis of the oil of *Heracleum persicum* L. (leaves and flowers). J Essent Oil Res 14(4): 295–7.
  12. Scheffer JJC, Hiltunen R, Aynehchi Y, von Schantz, M, Svendsen AB. 1984. Composition of essential oil of *Heracleum persicum* fruits. Planta Med 50(1): 56–60.
  13. George V, Chacko S, Sethuraman MG. 2001. Chemical composition of the essential oil from the rhizomes of *Heracleum candolleianum*. J Essent Oil Res 13(2): 80–1.
  14. Papageorgiou VP, Ochir G, Motl O, Argyriadou N, Dunkel H. 1985. Composition of the essential oil from *Heracleum dissectum*. J Nat Prod 48(5): 851–3.
  15. Işcan G, Demirci F, Kürkçüoğlu M, Kıvanç M, Başer KHC. 2003. The bioactive essential oils of *Heracleum sphondylium* L. subsp. *ternatum* (Velen.). Brummitt. Z Naturforsch 58c: 195–200
  16. Işcan G, Özek T, Özek G, Duran A, Başer KHC. 2004. Essential oils of tree species of *Heracleum* anticandidal activity. Chem Nat Comp 40(6): 544–7
  17. Tosun F, Akyüz Kızılay Ç, Erol K, Kiliç FS, Kürkçüoğlu M, Hüsnü Can Baserc K. 2008. Anticonvulsant activity of furanocoumarins and the essential oil obtained from the fruits of *Heracleum crenatifolium*. Food Chem 107(3): 990–3.
  18. John AJ, Karunakaran VP, George V, Sethuraman MG. 2007. Chemical composition of leaf and fruit oils of *Heracleum candolleianum*. J Essent Oil Res 19(4): 358–9.
  19. Bonjar GHS. 2004. Antibacterial screening of plants used in Iranian folkloric medicine. Fitoterapia 75(2): 231–5.
  20. Özek T, Özek G, Başer KHC, Duran A. 2005. Comparison of the essential oils of three endemic Turkish *Heracleum* species obtained by different isolation techniques. J Essent Oil Res 17(6): 605–10.
  21. Özek T, Demirci B, Başer KHC. 2002. Comparative study of the essential oils of *Heracleum sphondylium* ssp *ternatum* obtained by micro- and hydro-distillation methods. Chem Nat Comp 38(1): 48–50.
  21. Yáñez X, Pinzón ML, Solano F, Sánchez L R. 2002. Chemical composition of the essential oil of *Psidium caudatum* McVaugh. Molecules 7: 712–6.
  23. Adams RP. 1995. Identification of Essential Oil Components by Gas Chromatography-Mass Spectroscopy. In Allured Publishing: Carol Stream, Illinois, USA, p 475.
  24. Edwin HL, Albert B, William JJ, Jean HS. 1985. Manual of Clinical Microbiology. In 4 ed.; Washington, D. C., pp 978–7.
  25. Victor L. 1991. Antibiotics in laboratory medicine. In 3 ed.; Lippincott Williams & Wilkins: Maryland, USA, p 12.
  26. Jorgensen JH, Turnidge JD, Washington JA. 1999. Antibacterial susceptibility tests: Dilution and disk diffusion methods. In Manual of clinical microbiology, 7 ed.; Murry, PR.; Baron, EJ.; Pfaller, MA., Eds. ASM Press: Washington, D. C., pp 1526–43.
  27. Sriubolmas N Khampha W, Wiyakrutta S, Panphut W, Laowanapiban P, Plainkum P, et al. 1999. Microbial Sources of Bioactive Compounds and Enzymes Useful for Antibiotic Industry. In In Biotechnology for Sustainable Utilization of Biological Resources in the Tropics, Yoshida, T., et al., Eds. International Center for Biotechnology: Osaka,; Vol. 14, pp 165–82.
  28. Palanuvej C, Werawatganone P, Lipipun V, Ruangrunsi N. 2006. Chemical composition and antimicrobial activity against *Candida albicans*

- of essential oil from leaves of *Cinnamomum porrectum*. *J Health Res* 20(1): 69–76.
- 29.** Bakkali F, Averbeck S, Averbeck D, Idaomar M. 2008. Biological effects of essential oils – A review. *Food Chem. Toxicol* 46(2): 446–75.
- 30.** Wahab SMA, Aboutabl EA, El-Zalabani SM, Fouad HA, De Pooter HL, El-Fallaha B. 1987. The essential oil of olibanum. *Planta Med* 53(4): 382–4.
- 31.** Domokos J, Pálincás J, Héthelyi E, Korányi K, Perédi J. 2000. Examination on volatile and fatty oils of the seed of broas-leaved spignel (*Peucedanum cervaria* L.). *Acta Horti* 523: 97–104.
- 32.** Moshonas MG. 1971. Analysis of carbonyl flavor constituents from grapefruit oil. *J Agric Food Chem* 19(4): 769–0.
- 33.** Crowell PL, Elson CE, Bailey HH, Elegbede A, Haag JD, Gould MN. 1994. Human metabolism of the experimental cancer therapeutic agent *d*-limonene. *Canc Chemother Pharmacol* 35(1): 31–7.
- 34.** Sonboli A, Eftekhari F, Yousefzadi M, Kanani, MR. 2005. Antibacterial activity and chemical composition of the essential oil of *Grammosciadium platycarpum* Boiss. from Iran. *Z Naturforsch* 60c: 30–4.
- 35.** Sivropoulou A, Nikolaou C, Papanikolaou E, Kokkini S, Lanaras T, Arsenakis M. 1997. Antimicrobial, cytotoxic, and antiviral activities of *Salvia fruticosa* essential oil. *J Agric Food Chem* 45(8): 3197–201.
- 36.** Mikus J, Harkenthal M, Steverding D, Reichling J. 2000. *In vitro* effect of essential oils and isolated mono- and sesquiterpenes on *Leishmania major* and *Trypanosoma brucei*. *Planta Med* 66(4): 366–8.
- 37.** Shin S, Kim JH. 2004. Antifungal activities of essential oils from *Thymus quinquecostatus* and *T. magnus*. *Planta Med* 70(11): 1090–2.
- 38.** Reichling J, Suschke U, Schneelee J, Geiss HK. 2006. Antibacterial activity and irritation potential of selected essential oil components – Structure activity relationship. *Nat Prod Comm* 1 (11): 1003–12.
- 39.** Hoet S, Stévigny C, Hérent MF, Quetin-Leclercq J. 2006. Antitrypanosomal compounds from the leaf essential oil of *Strychnos spinosa*. *Planta Med* 72(5): 480–2.
- 40.** Franzios G, Mirotsoy M, HatziaPOSTOLOY E, Kral J, Scouras ZC, Mavragani-Tsipidou P. 1997. Insecticidal and genotoxic activities of mint essential oils. *J. Agric. Food Chem* 45(7): 2690–4.
- 41.** Santana-Rios G, Orner GA, Amantana A, Provost C, Wu S-Y, Dashwood RH. 2001. Potent antimutagenic activity of white tea in comparison with green tea in the *Salmonella* assay. *Mutat Res* 495(1–2): 61–74.
- 42.** Bruni R, et al. 2003. Chemical composition and biological activities of Isphingo essential oil, a traditional Ecuadorian spice from *Ocotea quixos* (Lam.) Kosterm. (Lauraceae) flower calices. *Food Chem* 85(3): 415–21.
- 43.** Sacchetti G, Maietti S, Muzzoli M, Scaglianti M, Manfredini S, Radice M, et al. 2005. Comparative evaluation of 11 essential oils of different origin as functional antioxidants, antiradicals and antimicrobials in food. *Food Chem* 91(4): 621–32.
- 44.** Werawatganone P, Palanuvej C, Lipipun V, Ruangrunsi N. 2006. Thep-ta-ro essential oil in solution and emulgel dosage forms. *J. Health Res* 20(1): 77–86.